





DEPARTMENT OF PARKS AND RECREATION OFFICE OF HISTORIC PRESERVATION

Julianne Polanco, State Historic Preservation Officer
1725 23rd Street, Suite 100, Sacramento, CA 95816-7100
Telephone: (916) 445-7000 FAX: (916) 445-7053
calshpo.ohp@parks.ca.gov www.ohp.parks.ca.gov

November 5, 2019

VIA EMAIL

In reply refer to: FHWA_2019_1015_002

Ms. Jody L. Brown, Office Chief Cultural Studies Office Division of Environmental Analysis Caltrans PO Box 942873, MS-27 Sacramento, CA 94273-0001

Subject: Determinations of Eligibility for the Proposed Last Chance Grade Phase 2B Geotechnical Investigations Project, Del Norte County, CA

Dear Ms. Brown:

Caltrans, as assigned by the Federal Highway Administration, is initiating consultation regarding the above project in accordance with Section 106 of the National Historic Preservation Act and implementing regulations codified at 36 CFR Part 800. As part of your documentation, Caltrans submitted a Historic Property Survey Report (HPSR), Historical Resources Evaluation Report, and an Archaeological Survey Report for the proposed project.

Caltrans District 1 proposes to develop a long-term solution to the instability and potential roadway failure at Last Chance Grade by constructing a new alignment. The existing alignment requiring replacement is located between PM 13.3 and 15.9 on US Highway 101 in Del Norte County, CA. There are currently several alternative alignments under consideration. Prior to completion of the design of alternative alignments, geotechnical studies are required to determine the depth and stability of soils and geological formations within the project area. A detailed description of the project and area of potential effect is located on pages 1-6 of the HPSR for the project.

As part of its identification efforts Caltrans determined that the Crescent City to Trinidad Wagon Road is eligible for the National Register of Historic Places (NRHP). Thirteen segments of the wagon road were identified for the purposes of this undertaking. Caltrans also determined that segments 1, 10, 11, and 13 retain historic integrity and are contributing elements of this resource. In addition Segments 2 through 9 and 12 do not retain historic integrity and are non-contributing elements to the wagon road.

FHWA_2019_1015_002

Ms. Brown November 5, 2019 Page 2 of 2

Caltrans has also determined that the Road Grade and Drainage Ditch, located adjacent to and up the slope from the 1930s alignment of the Redwood Highway (modern highway), is not eligible for the NRHP.

Pursuant to 36 CFR 800.4(a-c), Caltrans is requesting SHPO concurrence on the following:

- 1) Adequacy of the delineation of the APE
- 2) Adequacy of the identification effort
- 3) Adequacy of the evaluation of potential historic properties for eligibility to the NRHP.

I have reviewed the documentation furnished and have the following comments:

- 1) I have no comment with regards to the delineation of the APE.
- 2) With regards to the Crescent City to Trinity Wagon Road, I do not have enough information at this time to either agree or disagree with Caltrans' determination of eligibility for the NRHP. I recommend that Caltrans treat the property as eligible for the purposes of the project. For segments with compromised integrity, Caltrans can take these factors into account as part of the effects analysis.
- 3) I concur that the Road Grade and Drainage Ditch, located adjacent to and up the slope from the 1930s alignment of the Redwood Highway (modern highway), is not eligible for the NRHP.
- 4) I have no further comment regarding the identification effort.

If you have any questions, please contact Natalie Lindquist at (916) 445-7014 with e-mail at natalie.lindquist@parks.ca.gov.

Sincerely,

Julianne Polanco

State Historic Preservation Officer

DEPARTMENT OF TRANSPORTATION

DIVISION OF ENVIRONMENTAL ANALYSIS (MS 27) 1120 N STREET
P.O. BOX 942874
SACRAMENTO, CA 94274-0001
PHONE: (916) 654-3467
FAX: (916) 653-7757
TTY: (916) 653-4086



Make Conservation a California Way of Life

October 14, 2019

Ms. Julianne Polanco State Historic Preservation Officer 1725 23rd Street, Suite 100 Sacramento, CA 95816

Attn: Natalie Lindquist

Re: Determination of Eligibility for the Last Chance Grade Phase 2B Geotechnical Investigations Project, Del Norte County, California

Dear Ms. Polanco:

The Department of Transportation (Caltrans), as assigned by the Federal Highway Administration (FHWA), is initiating consultation with the State Historic Preservation Officer (SHPO) regarding the Last Chance Grade Phase 2B Geotechnical Investigations Project on US Highway 101, in Del Norte County, California.

The environmental review, consultation, and any other actions required by applicable Federal environmental laws for this project are being, or have been, carried out by Caltrans pursuant to 23 U.S.C. 327 and the Memorandum of Understanding dated December 23, 2016 and executed by FHWA and Caltrans. At the request of project stakeholder National Park Service, the Caltrans First Amended Section 106 Programmatic Agreement (January 2014) will not be utilized for this undertaking. Consultation will therefore occur under National Historic Preservation Act implementing regulations at 36 CFR § 800.

Caltrans, District 1, is proposing to develop a long-term solution to the instability and potential roadway failure at Last Chance Grade (LCG) by constructing a new alignment. The existing alignment requiring replacement is located between post miles (PM) 13.3 and 15.9 on US highway 101 in Del Norte County, California. There are currently several alternative alignments under consideration. Prior to the completion of the design of alternative alignments,

"Provide a safe, sustainable, integrated and efficient transportation system to enhance Califomia's economy and livability" Ms. Polanco Last Chance Grade Phase 2B Geotechnical Investigations October 14, 2019 Page 2

geotechnical studies are required to determine the depth and stability of soils and geological formations within the project area.

The proposed Phase 2B geotechnical investigations will include geotechnical borings at 16 locations and 14 seismic line surveys. The purpose of the exploration activities is to characterize sites in terms of depth to stable rock and rock properties, and to confirm the presence and activity of landslide features. Based on the materials encountered and the geotechnical design criteria desired, select soil samples will be obtained from borings for laboratory analysis. Pursuant to 36 CFR 800.3(g), Caltrans is requesting SHPO consultation on the appropriateness of the Area of Potential Effect (APE), the adequacy of historic property identification efforts, and determinations of eligibility for this undertaking.

In an effort to identify and evaluate historic properties, Caltrans has completed and attached the Historic Property Survey Report for the Last Chance Grade Phase 2B Geotechnical Investigation Project in Del Norte County, 01-DN-101 Post Mile 13.3-22.58 (HPSR), which contains information on the project and describes our efforts to identify and evaluate cultural resources. The HPSR can be referred to for additional information on the current undertaking. Project effects on historic properties will be discussed in a forthcoming Finding of Effects document.

As a result of archival research, record searches, archaeological surveys, and a Historical Resource Evaluation study, Caltrans has determined that two cultural resources within the APE require evaluation.

Caltrans has determined the following resource is **eligible** for listing in the National Register of Historic Places (NRHP) and is requesting your concurrence on eligibility:

• The Crescent City to Trinidad Wagon Road is the first north-south access route along the northern California coastline. Thirteen segments of the wagon road were identified and recorded for the purposes of this undertaking. Segments 1, 10, 11 and 13 retain historical integrity and are therefore contributing elements of this resource. Segments 2 through 9 and 12 do not retain historical integrity and are non-contributing elements to the wagon road.

"Provide a safe, sustainable, integrated and efficient transportation system to enhance California's economy and livability"

Ms. Polanco Last Chance Grade Phase 2B Geotechnical Investigations October 14, 2019 Page 3

Caltrans has determined the following resource is **ineligible** for listing in the National Register of Historic Places (NRHP) and is requesting your concurrence on eligibility:

 A Road Grade and Drainage Ditch is located adjacent to and up the slope from the 1930's alignment of the Redwood Highway (modern highway). The sshaped alignment appears to be an abandoned road associated with a drainage ditch constructed by the Division of Highways in the 1930's.

Caltrans' requests your concurrence regarding the adequacy of the following:

- Delineation of the Area of Potential Effects (APE) for the undertaking;
- Identification of potential historic properties located within the undertaking's APE
- Evaluation of potential historic properties for eligibility to the National Register of Historic Places (NRHP)

Thank you for your assistance in providing a safe, sustainable, integrated, and efficient transportation system to enhance California's economy and livability. If you require any additional information or have any questions or concerns please do not hesitate to contact me or Caltrans archaeologists Stacey Zolnoski at (707) 441-5855 or at Stacey. Zolnoski@dot.ca.gov, or Timothy Keefe at (707) 441-2022 or at Timothy. Keefe@dot.ca.gov, or Caltrans architectural historian Douglas Bright at (510) 286-5350 or at Douglas. Bright@dot.ca.gov.

Sincerely,

JÖDY L. BROWN

Office Chief

Cultural Studies Office

Enclosures:

Historic Property Survey Report for the Last Chance Grade Phase 2B Geotechnical Investigations, Del Norte County, California

> "Provide a safe, sustainable, integrated and efficient transportation system to enhance California's economy and livability"

Ms. Polanco Last Chance Grade Phase 2B Geotechnical Investigations October 14, 2019 Page 4

Copy: Timothy Keefe; Caltrans, District 1 Douglas Bright; Caltrans, District 4

> Alex Neeb, Caltrans, Cultural Studies Office Karin Grantham, Redwood National Park

Greg Collins; California State Parks, North Coast Redwoods District

Joe James and Rosie Clayburn; Yurok Tribe

Denise Padgette and Amanda O'Connell; Tolowa Dee-ni' Nation

Charlene Storr; Tolowa Nation

Dale Miller and Crista Stewart; Elk Valley Rancheria

Fawn Murphy; Resighini Rancheria

"Provide a safe, sustainable, integrated and efficient transportation system to enhance California's economy and livability"

OFFICE OF HISTORIC PRESERVATION DEPARTMENT OF PARKS AND RECREATION

1725 23rd Street, Suite 100 SACRAMENTO, CA 95816-7100 (916) 445-7000 Fax: (916) 445-7053 calshpo@parks.ca.gov www.ohp.parks.ca.gov

May 15, 2014

Reply To: FHWA_2014_0320_001

Steve Croteau, Chief Environmental Services Branch E1 Caltrans District 1 PO Box 3700 Eureka, CA 95502-3700

Re: Determination of Eligibility for the Proposed Storm Damage Permanent Restoration Project on Route 1, Del Norte County, CA

Dear Mr. Croteau:

You are consulting with me about the subject undertaking in accordance with the January 2014 First Amended Programmatic Agreement Among the Federal Highway Administration, the Advisory Council on Historic Preservation, the California State Historic Preservation Officer, and the California Department of Transportation Regarding Compliance with Section 106 of the National Historic Preservation Act, as it Pertains to the Administration of the Federal-Aid Highway Program in California (PA).

In 2001, Caltrans staff determined the Redwood Highway met the criteria for inclusion in the National Register of Historic Places (NRHP) pursuant to PRC 5024 under Criterion C as the design of a master landscape architect, an engineering achievement, and for its aesthetic qualities. In 2004, Caltrans sought and received concurrence from the SHPO for this determination.

Due to the passage of time and the discovery of new information, Caltrans reevaluated the Redwood Highway. Caltrans has now determined that the Redwood Highway is not eligible for the NRHP. Additionally as a state-owned resource, pursuant to PRC 5024(b), Caltrans has determined that the Redwood Highway does not meet California Historical Landmark eligibility criteria and should not be considered a historical resource for the purposes of CEQA. Based on my review of the submitted documentation, I concur.

Thank you for considering historic properties during project planning. If you have any questions, please contact Natalie Lindquist of my staff at (916) 445-7014 or email at natalie.lindquist@parks.ca.gov or Michelle Messinger at (916)445-7005 or e-mail at michelle.messinger@parks.ca.gov.

Sincerely,

Carol Roland-Nawi, Ph.D.

State Historic Preservation Officer

el Tokand Nair, Ph.D.

DEPARTMENT OF TRANSPORTATION

DISTRICT 1, P.O. BOX 3700 EUREKA, CA 95502-3700 PHONE (707) 441-5615 FAX (707) 441-5775 TTY 711



March 12, 2014

Carol Roland-Nawi, Ph.D. State Historic Preservation Officer 1725 23rd Street, Suite 100 Sacramento, CA 95816 US Route 101/DN/PM 15.1-15.14 Storm Damage Permanent Repair EA: 01-0B270/EFIS: 0112000111

Re: Historic Property Survey Report and Determination of Eligibility for the Storm Damage Permanent Restoration Project on Route 101, Del Norte County – Section 106 and PRC 5024 Compliance

Dear Dr. Roland-Nawi:

The California Department of Transportation (Caltrans) and Federal Highway Administration propose to repair US Route 101 within Del Norte Coast Redwoods State Park Bridge from post mile (PM) 15.1 to 15.14 in Del Norte County. The project is necessary because of substantial damage caused by winter storms in 2010 that caused a loss of a portion of the southbound traffic lane. A full project description can be found on pages one through two of the enclosed HPSR and a depiction of the Area of Potential Effects (APE) is attached to it (Attachment A). Caltrans is initiating consultation on this project in accordance with the January 2014 First Amended Programmatic Agreement Among the Federal Highway Administration, the Advisory Council on Historic Preservation, the California State Historic Preservation Officer, and the California Department of Transportation Regarding Compliance with Section 106 of the National Historic Preservation Act, as it Pertains to the Administration of the Federal-Aid Highway Program in California (PA).

Enclosed you will find a Historic Property Survey Report (HPSR) for the proposed undertaking with attached Historic Resources Evaluation Report (HRER). We are consulting with you at the present time under Stipulations VIII.C.6 of the PA, which requires that we seek your concurrence on Caltrans' determinations of eligibility for potential historic properties.

Caltrans is transmitting this study as a federal agency, following the provisions of 23 USC 326 and the *Memorandum of Understanding (MOU) between the Federal Highway Administration, California Division and the California Department of Transportation State Assumption of Responsibility for Categorical Exclusions*, which became effective on June 7, 2007, and was renewed on June 7, 2010. Please direct all future correspondence on this project to Caltrans.

Consultation and identification efforts for this project resulted in the identification of one previously recorded cultural resource within the APE:

 $"Caltrans\ improves\ mobility\ across\ California"$

Dr. Carol Roland-Nawi March 6, 2014 Page 2 of 2

> Redwood Highway through Del Norte Coast Redwoods State Park (Redwood Highway hereafter), PM 13.3-22.58

No other cultural resources are within the APE.

In 2001, Caltrans staff determined the Redwood Highway met the criteria for inclusion in the National Register of Historic Places (NRHP) pursuant to PRC 5024. It was determined to be a landscape district at the state level of significance under Criterion C as the design of a master landscape architect, an engineering achievement, and for its aesthetic qualities. In 2004, Caltrans sought and received formal concurrence from SHPO for this determination.

Due to the passage of time and the discovery of new information, Caltrans has reevaluated the Redwood Highway. Under Stipulation VIII.C.5 of the PA, Caltrans has concluded the previous determination is not valid and, pursuant to Stipulation VIII.C.6, is requesting your concurrence that the Redwood Highway is **not eligible** for listing in the NRHP. Additionally, as a state-owned resource, pursuant to PRC 5024(b), Caltrans requests your concurrence that it does not meet the California Historical Landmark eligibility criteria. Caltrans has also determined the Redwood Highway is not a historical resource for the purposes of CEQA.

We look forward to receiving your response within 30 days of receipt of this submittal in accordance with Stipulation VIII.C.6.a of the PA. Please contact Chris Kuzak, North Region Architectural Historian, at (530) 741-4017, if you have any questions regarding the documentation enclosed with this letter.

Sincerely,

Steven Croteau, Chief

Environmental Services Branch E1

teve Crotean

Enclosure

cc: Michelle Messenger, OHP

"Caltrans improves mobility across California"



Appendix F. Biological Surveys – Species, Personnel, and Dates



Survey Date(s)	Survey Type	Personnel	Organization
March 26, 2019	Special-Status Animal Habitat Assessment	Christine Hamilton (biologist)	Caltrans
April 15, 2019	Botanical Survey	Jeff Barrett (botanist), Jon Lee (botanist)	Caltrans
April 16, 2019	Botanical Survey	Jeff Barrett (botanist), Amanda Lee (project coordinator), Christine Hamilton (biologist), Jon Lee (botanist)	Caltrans
April 17, 2019	Botanical Survey	Jeff Barrett (botanist), Amanda Lee (project coordinator), Jacob Hillard (botanist), Jon Lee (botanist)	Caltrans
April 18, 2019	Botanical Survey	Jeff Barrett (botanist), Felicia Zimmerman (project coordinator), Christine Hamilton (biologist), Jon Lee (botanist), Jacob Hillard (botanist)	Caltrans
April 19, 2019	Botanical Survey	Jeff Barrett (botanist), Amanda Lee (project coordinator), Christine Hamilton (biologist), Jacob Hillard (botanist)	Caltrans
April 26, 2019	Botanical Survey	Jeff Barrett (botanist), Amanda Lee (project coordinator), Felicia Zimmerman (project coordinator), Kellie Eldridge (botanist)	Caltrans
May 1, 2019	Botanical Survey	Jeff Barrett (botanist), Christine Hamilton (biologist), Jon Lee (botanist), Jacob Hillard (botanist)	Caltrans
May 2, 2019	Botanical Survey	Jeff Barrett (botanist), Amanda Lee (project coordinator), Jon Lee (botanist), Felicia Zimmerman,	Caltrans
May 3, 2019	Botanical Survey	Jeff Barrett (botanist), Amanda Lee (project coordinator), Christine Hamilton (biologist), Felicia Zimmerman (project coordinator)	Caltrans
May 15, 2019	Botanical Survey	Jeff Barrett (botanist), Christine Hamilton (biologist), Amanda Lee (project coordinator), Jon Lee (botanist)	Caltrans
May 16, 2019	Botanical Survey	Jeff Barrett (botanist), Christine Hamilton (biologist), Ali Thiel (biologist), Jon Lee (botanist)	Caltrans
May 22-24, 2019	Aquatic Resources Delineation	Jordan Mayor (wetland ecologist), Margaret Widdowson (botanist/wetland ecologist)	ICF
June 11-12, 2019	Botanical Survey	Jeff Barrett (botanist), Jon Lee (botanist), Jacob Hillard (botanist), Christine Hamilton (biologist)	Caltrans

Survey Date(s)	Survey Type	Personnel	Organization
June 13, 2019	Botanical Survey	Jeff Barrett (botanist), Jon Lee (botanist), Jacob Hillard (botanist),	Caltrans
June 19, 2019	Botanical Survey	Jeff Barrett (botanist), Jacob Hillard (botanist), Jon Lee (botanist), Kellie Eldridge (botanist)	Caltrans
June 23-26, 2019	Aquatic Resources Delineation	Jordan Mayor (wetland ecologist), Zach Larson (biologist)	ICF
July 17-18, 2019	Aquatic Resources Delineation	Jordan Mayor (wetland ecologist), Margaret Widdowson (botanist/wetland ecologist)	ICF
July 23-24, 2019	Botanical Survey	Jeff Barrett (botanist), Jon Lee (botanist), Jacob Hillard (botanist), Jeremy Pohlman (biologist)	Caltrans
July 25, 2019	Botanical Survey	Jeff Barrett (botanist), Jon Lee (botanist), Jeremy Pohlman (biologist), Felicia Zimmerman (project coordinator)	Caltrans
July 29, 2019	Botanical Survey	Jeff Barrett (botanist), Ali Thiel (biologist)	Caltrans
July 30, 2019	Botanical Survey	Jeff Barrett (botanist), Ali Thiel (biologist), Stacey Zolnoski (archaeologist)	Caltrans
July 31, 2019	Botanical Survey	Jeff Barrett (botanist), Ali Thiel (biologist), Kellie Eldridge (botanist)	Caltrans
October 8, 2019	Aquatic Resources Assessment	Robert Meade (senior biologist), Jeff Barrett (botanist), Kellie Eldridge (botanist)	Caltrans
October 25, 2019	Aquatic Resources Delineation	Jeff Barrett (botanist), Kellie Eldridge (botanist), Alexandra Thiel (biologist)	Caltrans

Appendix G. Special-Status Plant Species with the Potential to Occur in the Project Vicinity



Common Name/ Scientific Name	Status ^a	Habitat Requirements and Blooming Period	Habitat Present/ Absent ^b	Rationale
Sea-watch/ Angelica lucida	CRPR 4.2	Coastal bluff scrub, coastal dunes, coastal scrub, marshes and swamps (coastal salt). Occurs from 0 to 150 meters in elevation; blooms May–September.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
False gray horsehair lichen/ Bryoria pseudocapillaris	CRPR 3.2	Coastal dunes (San Luis Obispo County), North Coast coniferous forest (immediate coast). Occurs from 0 to 90 meters in elevation.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Twisted horsehair lichen/ Bryoria spiralifera	CRPR 4.2	North Coast coniferous forest. Occurs from 0 to 30 meters in elevation.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Bolander's reed grass/ Calamagrostis bolanderi	CRPR 4.2	Bogs and fens, broadleafed upland forest, closed-cone coniferous forest, coastal scrub, meadows and seeps (mesic), marshes and swamps (freshwater), North Coast coniferous forest. Occurs from 0 to 455 meters in elevation; blooms May–August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Thurber's reed grass/ Calamagrostis crassiglumis	CRPR 2B.1	Coastal scrub, freshwater marsh, marsh and swamp, wetland. Occurs from 0 to 50 meters in elevation; blooms May–August.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Leafy reed grass/ Calamagrostis foliosa	SR, CRPR 4.2	Coastal bluff scrub, North Coast coniferous forest. Occurs from 0 to 1,220 meters in elevation; blooms May–September.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Spiral-spored gilded- head pin lichen/ Calicium adspersum	CRPR 2B.1	Lower montane coniferous forest, North Coast coniferous forest.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Seaside bittercress/ Cardamine angulata	CRPR 2B.1	Lower montane coniferous forest, North Coast coniferous forest, wetlands. Occurs from five to 915 meters in elevation; blooms January–July.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Northern clustered sedge/ Carex arcta	CRPR 2B.2	Bogs and fens, North Coast coniferous forest, Wetlands. Occurs from 60 to 1,400 meters in elevation; blooms June–September.	Р	Low: Small amount of suitable habitat is present with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).

Common Name/ Scientific Name	Status ^a	Habitat Requirements and Blooming Period	Habitat Present/ Absent ^b	Rationale
Buxbaum's sedge Carex buxbaumii	CRPR 4.2	Bogs and fens, meadows and seeps, marshes and swamps.		Low: Minor amount of suitable habitat present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Bristle-stalked sedge Carex leptalea	CRPR 2B.2	Bogs and fens, freshwater marsh, marshes and swamps, meadows and seeps, wetlands.	Р	Low: Minor amount of suitable habitat present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Northern meadow sedge Carex praticola	CRPR 2B.2	Meadows and seeps, wetlands.	Р	Low: Minor amount of suitable habitat present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Green yellow sedge Carex viridula ssp. viridula	CRPR 2B.3	North Coast coniferous forests; bogs and fens, marshes and swamps, wetlands.	Р	Low: Minor amount of suitable habitat at present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Nuttall's saxifrage/ Cascadia nuttallii	CRPR 2B.1	North Coast coniferous forest. Occurs from 35 to 80 meters in elevation; blooms in May.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Johnny-nip/ Castilleja ambigua ssp. ambigua	CRPR 4.2	Coastal bluff scrub, coastal prairie, coastal scrub, marshes and swamps, valley and foothill grassland, vernal pool margins. Occurs from 0 to 435 meters in elevation; blooms March–August.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Oregon coast paintbrush Castilleja litoralis	CRPR 2B2	Coastal bluff scrub, coastal dunes, and coastal scrub.	Р	Moderate: suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County)
Oregon coast paintbrush/ Castilleja litoralis	CRPR 2B.2	Coastal bluff scrub, coastal dunes, coastal scrub. Occurs from five to 255 meters in elevation; blooms June–July.		Moderate: suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County)
Pacific golden- saxifrage/ Chrysosplenium glechomifolium	CRPR 4.3	North Coast coniferous forest, riparian forest. Occurs from 10 to 455 meters in elevation.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.

Common Name/ Scientific Name	Status ^a	Habitat Requirements and Blooming Period	Habitat Present/ Absent ^b	Rationale
Oregon goldthread/ Coptis laciniata	CRPR 4.2	Meadows and seeps, North Coast coniferous forest (streambanks). Occurs from 0 to 1,000 meters in elevation; blooms February– November.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Naked flag moss/ Discelium nudum	CRPR 2B.2	Coastal bluff scrub (soil, on clay banks). Occurs from 10 to 50 meters in elevation.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Black crowberry/ Empetrum nigrum	CRPR 2B.2	Coastal bluff scrub, coastal prairie. Occurs from 3 to 200 meters in elevation; blooms April–June.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Del Norte buckwheat/ Eriogonum nudum var. paralinum	CRPR 2B.2	Coastal bluff scrub, coastal prairie. Occurs from five to 80 meters in elevation; blooms June–September.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Coast fawn lily/ Erythronium revolutum	CRPR 2B.2	Bogs and fens, broadleafed upland forest, North Coast coniferous forest. Occurs from 0 to 1,600 meters in elevation; blooms March August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Minute pocket moss/ Fissidens pauperculus	CRPR 1B.2	North Coast coniferous forest (damp coastal soil). Occurs from 10 to 1,024 meters in elevation.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Pacific gilia/ Gilia capitata ssp. pacifica	CRPR 1B.2	Coastal bluff scrub, chaparral (openings), coastal prairie, valley and foothill grassland. Occurs from five to 1,665 meters in elevation; blooms April–August.	Р	Low: Minor amount of suitable habitat is present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County)
Harlequin lotus/ Hosackia gracilis CRPR 4.2		Broadleafed upland forest, coastal bluff scrub, closed-cone coniferous forest, Cismontane woodland, coastal prairie, coastal scrub, meadows and seeps, marshes and swamps, North Coast coniferous forest, valley and foothill grassland (wetlands, roadsides). Occurs from 0 to 700 meters in elevation; blooms March–July.	Р	Low: Minor amount of suitable habitat is present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County)

Common Name/ Scientific Name	Status ^a	Habitat Requirements and Blooming Period	Habitat Present/ Absent ^b	Rationale
California globe mallow/ Iliamna latibracteata	CRPR 1B.2	Chaparral (montane), lower montane coniferous forest, North Coast coniferous forest (mesic), riparian scrub (streambanks). Occurs from 60 to 2,000 meters in elevation; blooms June–August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Thompson's iris/ Iris thompsonii	CRPR 4.3	Lower montane coniferous forest, North Coast coniferous forest. Occurs from 90 to 600 meters in elevation; blooms March–August.	Р	Low: Small amount of suitable habitat is present, with a few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Small groundcone/ Kopsiopsis hookeri	CRPR 2B.3	North Coast coniferous forest. Occurs from 90 to 1,435 meters in elevation; blooms April–August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Marsh pea/ Lathyrus palustris	CRPR 2B.2	Bogs and fens, coastal prairie, coastal scrub, lower montane coniferous forest, marshes and swamps, North Coast coniferous forest. Occurs from 1 to 140 meters in elevation; blooms March–August.	Р	Low: Minor amount of suitable habitat is present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County)
Kellogg's lily/ Lilium kelloggii	CRPR 4.3	Lower montane coniferous forest, North Coast coniferous forest. Occurs from 3 to 1,300 meters in elevation; blooms May–August.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Western lily/ Lilium occidentale	FE/SE/ CRPR 1B.1	Bogs and fens, coastal bluff scrub, coastal prairie, coastal scrub, marshes and swamps (freshwater), North Coast coniferous forest (openings). Occurs from 2 to 185 meters in elevation; blooms June–July.	Р	Low: Minor amount of suitable habitat is present, or few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County)
Vollmer's lily/ Lilium pardalinum ssp. vollmeri	CRPR 4.3	Bogs and fens, meadows and seeps (mesic). Occurs from 30 to 1,680 meters in elevation; blooms June–August.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Heart-leaved twayblade/ Listera cordata	CRPR 4.2	Bogs and fens, lower montane coniferous forest, North Coast coniferous forest. Occurs from five to 1,370 meters in elevation; blooms February–July.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.

Common Name/ Scientific Name	Statusª	Habitat Requirements and Blooming Period Habitat Present/ Absent ^b		Rationale
Coast Range lomatium/ Lomatium martindalei	CRPR 2B.3	Coastal bluff scrub, lower montane coniferous forest, meadows and seeps. Occurs from 240 to 3,000 meters in elevation; blooms May–August.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Inundated bog club- moss Lycopodiella inundata	CRPR 2B.2	Lower montane coniferous forests; bogs and fens, marshes and swamps.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Running-pine/ Lycopodium clavatum	CRPR 4.1	Lower montane coniferous forest (mesic), marshes and swamps, North Coast coniferous forest (mesic). Occurs from 45 to 1,225 meters in elevation; blooms June–September.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Leafy-stemmed miterwort/ Mitellastra caulescens	CRPR 4.2	Broadleafed upland forest, lower montane coniferous forest, meadows and seeps, North Coast coniferous forest. Occurs from five to 1,700 meters in elevation; blooms March–October.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Woodnymph/ Moneses uniflora	CRPR 2B.2	Broadleafed upland forest, North Coast coniferous forest. Occurs from 50-1,100 meters in elevation; blooms May–August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Ghost-pipe/ Monotropa uniflora	CRPR 2B.2	Broadleafed upland forest, North Coast coniferous forest. Occurs from 10 to 855 meters in elevation; blooms June-September.		High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Howell's montia/ Montia howellii	CRPR 2B.2	Meadows and seeps, North Coast coniferous forest, vernal pools. Occurs from 0 to 1,215 meters in elevation; blooms January–May.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Wolf's evening- primrose/ Oenothera wolfii	CRPR 1B.1	Coastal bluff scrub, coastal dunes, coastal prairie, lower montane coniferous forest. Occurs from 0- to 800 meters in elevation; blooms May–October.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Suksdorf's wood- sorrel/ Oxalis suksdorfii	CRPR 4.3	Broadleafed upland forest, North Coast coniferous forest. Occurs from 15 to 700 meters in elevation; blooms May–August.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.

Common Name/ Scientific Name	Status ^a	Habitat Requirements and Blooming Period Habitat Present/ Absent ^b		Rationale
Seacoast ragwort/ Packera bolanderi var. bolanderi	CRPR 2B.2	Coastal scrub, North Coast coniferous forest. Occurs from 30 to 915 meters in elevation; blooms January–August.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
White-flowered rein orchid/ Piperia candida	CRPR 1B.2	Broadleafed upland forest, lower montane coniferous forest, North Coast coniferous forest. Occurs from 20 to 1,615 meters in elevation; blooms March—September.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
California pinefoot/ Pityopus californicus	CRPR 4.2	Broadleafed upland forest, lower montane coniferous forest, North Coast coniferous forest, upper montane coniferous forest. Occurs from 15 to 2,225 meters in elevation; blooms March–August.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Nodding semaphore grass/ Pleuropogon refractus	CRPR 4.2	Lower montane coniferous forest, meadows and seeps, North Coast coniferous forest, riparian forest. Occurs from 0 to 1,600 meters in elevation; blooms March–August.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Oregon polemonium/ Polemonium carneum	CRPR 2B.2	Coastal prairie, coastal scrub, lower montane coniferous forest. Occurs from 0 to 1,830 meters in elevation; blooms April–September.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Angel's hair lichen/ Ramalina thrausta	CRPR 2B.1	North Coast coniferous forest. Occurs from 75 to 430 meters in elevation.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Trailing black currant/ Ribes laxiflorum	CRPR 4.3	North Coast coniferous forest. Occurs from five to 1,395 meters in elevation; blooms March–August.		High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Great burnet/ Sanguisorba officinalis	CRPR 2B.2	Bogs and fens, broadleaved upland forest, marshes and swamps, meadows and seeps, North Coast coniferous forest, riparian forest, ultramafic, wetlands. Occurs from 60 to 1,400 meters in elevation; blooms July-October.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).

Common Name/ Scientific Name	Status ^a	Habitat Requirements and Blooming Period Absen		Rationale
Maple-leaved checkerbloom/ Sidalcea malachroides	CRPR 4.2	Broadleafed upland forest, coastal prairie, coastal scrub, North Coast coniferous forest, riparian woodland. Occurs from 0 to 765 meters in elevation; blooms March–August.	Р	High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Siskiyou checkerbloom/ Sidalcea malviflora ssp. patula	CRPR 1B.2	Coastal bluff scrub, coastal prairie, North Coast coniferous forest. Occurs from five to 1,255 meters in elevation; blooms April–August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Coast checkerbloom/ Sidalcea oregana ssp. eximia	CRPR 1B.2	Lower montane coniferous forest, meadows and seeps, North Coast coniferous forest, wetlands. Occurs from five to 1,805 meters in elevation; blooms June–August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Scouler's catchfly/ Silene scouleri ssp. scouleri	CRPR 2B.2	Coastal bluff scrub, coastal prairie, valley and foothill grassland. Occurs from 0 to 600 meters in elevation; blooms March–September.	Р	Low: Small amount of suitable habitat is present, with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Trifoliate laceflower/ Tiarella trifoliata var. trifoliata	CRPR 3.2	Lower montane coniferous forest, North Coast coniferous forest (edges, moist shady banks, streambanks). Occurs from 170 to 1,500 meters in elevation; blooms May–August.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Cylindrical trichodon/ Trichodon cylindricus	CRPR 2B.2	Broadleafed upland forest, meadows and seeps, upper montane coniferous forest. Occurs from 50 to 2,002 meters in elevation.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Coastal triquetrella/ Triquetrella californica	CRPR 1B.2	Coastal bluff scrub, coastal scrub. Occurs from 10 to 100 meters in elevation.	Р	Moderate: Suitable habitat is present, and occurrences are recorded from the region (coastal Del Norte County and northern coastal Humboldt County).
Methuselah's beard lichen/ Usnea longissima	CRPR 4.2	Broadleafed upland forest, North Coast coniferous forest. Occurs from 45 to 1,465 meters in elevation.		High: Suitable habitat is present, and occurrences are recorded within five miles of ESL.
Alpine marsh violet/ Viola palustris	CRPR 2B.2	elevation. Bogs and fens, coastal scrub, wetlands. Occurs from 0 to 150 meters in elevation; blooms March— August.		Low: Small amount of suitable habitat is present with few to no occurrences recorded from the region (coastal Del Norte County and northern coastal Humboldt County).

a Status:

FE Federal Endangered. SE = State Endar SR = State Rare. State Endangered.

California Rare Plant Rank (CRPR)

- 1B
- = Rank 1B species: rare, threatened, or endangered in California and elsewhere.
 = Rank 2B species: rare, threatened, or endangered in California but more common elsewhere. 2B
- Review List: Plants about which more information is needed. 3
- Watch List: Plants of limited distribution. 4

Threat rank extensions:

- Seriously threatened in California (over 80% of occurrences threatened/high degree and immediacy of threat).
 Moderately threatened in California (20%-80% occurrences threatened/ moderate degree and immediacy of threat).
- .3 Not very threatened in California (less than 20% of occurrences threatened / low degree and immediacy of threat or no current threats).

Habitat Present/Absent:

- = Present (species is present).
- HP Habitat present (habitat is, or may be present).
- CH = Critical habitat present.

Appendix H. Special-Status Wildlife Species with Potential to Occur in the Project Vicinity



Common	Scientific Name	Status	Habitat Requirements	Habitat Present/	Rationale
Name	Scientific Name	Otatus	Habitat Requirements	Absent ^b	Rationale
			Amphibians		
Pacific (coastal) tailed-frog	Ascaphus truei	SSC	Typically found in cold, clear, perennial rocky streams in wet forests, but may also utilize intermittent creeks. This species is not known to use ponds or lakes. This species may occur in uplands during precipitation events.	Р	Intermittent streams mapped at or near SL-16, SL-21, B-16, B-20, B-35, and B-36 may provide aquatic habitat during winter and spring months when water is present.
Northern red- legged frog	Rana aurora	SSC	Humid forest, woodlands, grasslands, and stream sides in northwestern California, usually near dense riparian vegetation.	Р	Intermittent streams mapped at or near SL-16, SL-21, B-16, B-20, B-35, and B-36 may provide aquatic habitat during winter and spring months when water is present. May occur in forested uplands during the wet season near these same locations.
Foothill yellow-legged frog	Rana boylii	SCT	Species occurs throughout the North and South Coast Ranges, south to the Transverse Range, across northern California to the west slope of the Cascade Range, and south through the foothills of the Sierra Nevada. Occurs up to 6,000 feet in the northern Sierra Nevada. Inhabits forest streams and rivers with sunny, sandy, and rocky banks, with deep pools, and shallow riffles. Occurs in both perennial and intermittent streams.	Р	Intermittent streams mapped at or near SL-16, SL-21, B-16, B-20, B-35, and B-36 may provide aquatic habitat during winter and spring months when water is present.
Southern torrent salamander	Rhyacotriton variegatus	SSC	Occurs along the coast in cold and well-shaded perennial streams and seeps in hardwood and coniferous forests. Eggs are laid in flowing water and adults are typically found among moss-covered pebbles and rocks within or adjacent to flowing water.	Р	Intermittent streams mapped at or near SL-16, SL-21, B-16, B-20, B-35, and B-36 may provide aquatic habitat during winter and spring months when water is present.
	T		Reptiles	T	
Green sea turtle, East Pacific Distinct Population Segment (DPS)	Chelonia mydas	FT	Does not nest on beaches of northern California (NMFS and USFWS 2007); may occur in open water habitat off the coast line of Del Norte County.	A	No habitat in BSA.
Leatherback sea turtle	Dermochelys coriacea	FE	No known nesting sites on the coast of California (NMFS 2019); may occur in open water habitat off the coast of Del Norte County.	А	No habitat in BSA.
Western pond turtle	Emys marmorata	SSC	Occurs throughout California west of the Sierra-Cascade crest; found from sea level to 6,000 feet; does not occur in desert regions except along the Mojave River and its tributaries; occupies ponds, marshes, rivers, streams, and irrigation canals with muddy or rocky bottoms.	А	No habitat in BSA.
Olive ridley sea turtle	Lepidochelys olivacea	FE	Mainly pelagic in tropical/temperate regions of Pacific, South Atlantic, and Indian Oceans but has been known to inhabit coastal areas, including bays and estuaries.	А	No habitat in BSA.
			Birds		· · · · · · · · · · · · · · · · · · ·

Common Name	Scientific Name	Status ^a	Habitat Requirements	Habitat Present/ Absent ^b	Rationale
Marbled murrelet	Brachyramphus marmoratus	FT/SE	Nests in old-growth coniferous forests, up to 6 miles inland, often in Douglas-fir. Feeds near-shore; nests inland along coast from Eureka to Oregon border and from Half Moon Bay to Santa Cruz.	P, CH	May occur in mature forested habitat within the BSA. Known from project vicinity. Some project locations within critical habitat.
Western snowy plover	Charadrius alexandrinus nivosus	FT/SSC	Coastal beaches, sandy areas near estuaries, salt ponds, river mouths, and levees along inland salt ponds. Nests on the ground, mainly in the open in sandy areas. The BSA is not within critical habitat for this species.	А	No foraging or nesting habitat in BSA.
Northern harrier	Circus hudsonius	SSC	Nests on the ground among herbaceous vegetation, such as grasses or cattails; forages in grasslands, agricultural fields, and marshes. Breeding range encompasses much of lowland California; winter range expands to include the remaining lowland areas.	А	No foraging or nesting habitat in BSA.
Yellow-billed cuckoo	Coccyzus americanus	FT/SE	Prefers riparian woodlands composed of various compositions with a dense understory along slowmoving watercourses. Typically requires 25–99 acres of expansive riparian habitat for breeding.	А	No foraging or nesting habitat in BSA.
Black swift	Cypseloides niger	SSC	Colonial breeder on cliffs behind or adjacent to waterfalls and sea-bluffs above the surf; forages widely.	А	No foraging or nesting habitat in BSA.
White-tailed kite	Elanus leucurus	SFP	Forages in grasslands, meadows, or marshes. Nests in woodlands and mature riparian habitat types.	Α	No foraging or nesting habitat in BSA.
Tufted puffin	Fratercula cirrhata	SSC	Feeds in the ocean; nests along the coast on islands, islets, and mainland cliffs.	А	No foraging or nesting habitat in BSA.
Bald eagle	Haliaeetus leucocephalus	FD/SE/S FP	Nests in large, old-growth, or dominant live trees. Nests typically located 50–200 feet above ground. Forages primarily in large inland fishbearing waters with adjacent large trees or snags, and occasionally in uplands with abundant rabbits, other small mammals, or carrion.	А	Unlikely to occur in BSA; no foraging habitat and marginal nesting habitat.
Fork-tailed storm-petrel	Oceanodroma furcata	ssc	Colonial nester on islands. Forages over the open ocean.	А	No foraging or nesting habitat in BSA.
Short-tailed albatross	Phoebastria (=Diomedea) albatrus	FE/SSC	Pelagic species. Does not breed or nest in California; only known breeding sites in Western Pacific Ocean islands. During nonbreeding season, may occur in northern California along the shelf break of the continental shelf (USFWS 2008).	А	No foraging or nesting habitat in BSA.
Northern spotted owl	Strix occidentalis caurina	FT/ST	Mature old growth forests, conifers, wooded canyons; BSA is not within critical habitat for this species. Coniferous forests with a multilayered, multispecies canopy with moderate to high canopy closure; large snags (standing dead trees); an abundance of large, dead wood on the ground; and open space within and below the upper canopy to fly.	Р	May occur in mature forested habitat within BSA. Known from project vicinity.

Common Name	Scientific Name	Status ^a	Habitat Requirements	Habitat Present/ Absent ^b	Rationale
Little willow flycatcher	Empidonax Traillii brewsteri	SE	Prefers mountain meadows and riparian habitats. Nests near the edges of vegetation clumps and near streams in mountain meadows and riparian habitats.	А	No foraging or nesting habitat in BSA.
Long-eared owl	Asio otus	SSC	Require adjacent open land, productive of mice and the presence of old nests of crows, hawks, or magpies for breeding.	А	No foraging or nesting habitat in BSA.
Olive-sided flycatcher	Contopus cooperi	SSC	Summer resident; Most numerous in montane conifer forests where tall trees overlook canyons, meadows, lakes or other open terrain.	А	Species is a summer resident to the area and would not be present during project activities.
Peregrine falcon	Falco peregrinus anatum	SFP	Near wetlands, lakes, rivers, or other water; on cliffs, banks, dunes, mounds; also, humanmade structures. Nest consists of a scrape or a depression or ledge in an open site.	А	No foraging or nesting habitat in BSA.
Purple martin	Progne subis	SSC	Summer resident; Inhabits woodlands, low elevation coniferous forest of Douglas-fir, ponderosa pine, and Monterey pine.	А	Species is a summer resident to the area and would not be present during project activities.
Vaux's swift	Chaetura vauxi	SSC	Summer resident; Redwood, Douglas-fir, & other coniferous forests. Nests in large hollow trees & snags. Often nests in flocks.	А	Species is a summer resident to the area and would not be present during project activities.
Yellow- breasted chat	Icteria virens	SSC	Summer resident; inhabits riparian thickets of willow and other brushy tangles near watercourses. Nests in low, dense riparian habitat, consisting of willow, blackberry, wild grape; forages and nests within 10 feet of ground.	А	No suitable habitat in BSA.
Yellow warbler	Setophaga petechial	SSC	Prefers riparian plant associations near water. Frequently found nesting and foraging in willow shrubs and thickets, and in other riparian plants including cottonwoods, sycamores, ash, and alders.	А	No suitable habitat in BSA.
	T		Mammals		
Ring-tailed cat	Bassariscus astutus	SFP	Coniferous forests and riparian areas in California. Inhabit rock crevices, tree hollows, and cliffs.	Р	Suitable habitat in BSA.
Sonoma tree vole	Arborimus pomo	SSC	Occurs in old-growth and other forests, mainly Douglas-fir, redwood, and montane hardwood-conifer habitats.	Р	Suitable habitat in BSA.
White-footed vole	A. albipes	SSC	In California, only known from Humboldt and Del Norte Counties. Found in coastal forests dominated by redwood, Douglas-fir, and also occurs in riparian forest cover types. Occupies habitat near small streams with dense alder and deciduous trees and shrubs.	Р	Suitable habitat in BSA.
Sei whale	Balaenoptera borealis	FE	Worldwide cosmopolitan distribution in subtropical, temperate, and subpolar waters; usually observed in deeper waters of oceanic areas far from coastline.	А	No suitable habitat in BSA.

Common Name	Scientific Name	Status ^a	Habitat Requirements	Habitat Present/ Absent ^b	Rationale
Blue whale	Balaenoptera musculus	FE	Worldwide, often near the edges of physical features where krill tend to concentrate.	А	No suitable habitat in BSA.
Fin whale	Balaenoptera physalus	FE	Deep, offshore waters of all major oceans; less common in the tropics.	Α	No suitable habitat in BSA.
Townsend's big-eared bat	Corynorhinus townsendii	SSC	Primarily roosts in caves and cave- like roosting habitat, such as tunnels and mines. Very sensitive to disturbances and may abandon a roost after one onsite visit. Reported to use buildings in the northern and coastal portions of range. Also reported to use bridges (typically the cavernous spaces underneath) and hollow trees as roost sites. In California, occurs in inland deserts, moist cool redwood forests, oak woodlands of the inner Coast Ranges and Sierra Nevada foothills, and lower- to mid-elevation mixed coniferous forests.	Р	Mature redwood trees with cavities may provide suitable roosting sites; may forage throughout the BSA.
North Pacific right whale	Eubalaena japonica	FE	North Pacific Ocean; seasonally migratory; colder waters for feeding, migrating to warmer waters for breeding and calving; may move far out to sea during feeding seasons but gives birth in coastal areas.	А	No suitable habitat present in BSA.
Humboldt marten	Martes caurina humboldtensis	FPT/SE	Coastal old-growth forests and serpentine areas.	Р	Suitable habitat in BSA. Known from project vicinity.
Humpback whale	Megaptera novaeangliae	FT	All major oceans; central California population migrates from winter calving and mating areas off Mexico to summer and fall feeding areas off coastal California. Humpback whales occur from late April to early December.	А	No suitable habitat present in BSA.
Southern resident killer whale	Orcinus orca	FE	All oceans; most abundant in colder waters but also occurs in temperate water; presence and occurrence common but unpredictable in coastal California.	А	No suitable habitat present in BSA.
Fisher, West Coast DPS / Northern California Evolutionary Significant Unit (ESU)	Pekania pennanti	FPT/SSC	Requires forest with dense canopy and a complex structure that includes lots of down wood, moderate shrub cover, dead trees, and intermixed hardwood trees. Relies on dens created by large trees, snags, logs, rock piles, and root burrows.	Р	Suitable habitat in BSA. Known from project vicinity.
Sperm whale	Physeter catodon [=microcephalus]	FE	Open ocean far from land and uncommon in waters less than 984 feet (300 meters) deep; live at surface of the ocean but dive deep to catch giant squid.	А	No suitable habitat present in BSA.
Fish					
Green sturgeon, southern DPS	Acipenser medirostris	FT	Marine and estuarine environments, Sacramento River; San Francisco Bay-Delta, Humboldt Bay, offshore waters to 110 meters (360 feet) from Monterey Bay to the United States- Canada border.	Α	No suitable habitat present in BSA.

Common Name	Scientific Name	Status ^a	Habitat Requirements	Habitat Present/ Absent ^b	Rationale
Tidewater goby	Eucyclogobius newberryi	FE/SSC	Inhabits lagoons and estuaries with still or slow-moving water less than 3 feet deep. Salinity levels typically less than 12 parts per thousand, although they have been found in water with salinity from 0 to 42 parts per thousand, temperatures from 8 to 25degrees Celsius (°C) (46–77°F). Typically occurs over a sandy or mixed sandy/silty bottom with sparse vegetation.	А	No suitable habitat present in BSA.
Coho salmon, Southern Oregon/North ern California Coast ESU	Oncorhynchus kisutch	FT/ST	Found in perennial streams with water temperatures of 12–14°C. Not commonly found in streams where summer temperatures exceed 22–25°C. Requires deep pools, riffles, and runs with adequate canopy cover.	А	No suitable habitat present in BSA.
Chinook salmon, California Coastal ESU	Oncorhynchus tshawytscha	FT	Spends between 1 and 5 years in the ocean before returning to natal rivers to spawn. Typically enters freshwater river systems after large winter storm events. Spawns between October and December in the upper mainstems of rivers and the lower reaches of coastal creeks (Moyle et al. 2008) composed of a mixture of small cobble and large gravel.	А	No suitable habitat present in BSA.
Steelhead, Northern California DPS	O. mykiss irideus	FT, SCE	Live as adults in ocean habitats and migrate into rivers and streams to spawn (Moyle 2002). Steelhead spawn in gravel and small cobble substrates usually associated with riffle-and-run habitat types in coldwater streams.	А	No suitable habitat present in BSA.
Coast cutthroat trout	O. clarkii	SSC	Occupies coastal streams with some populations migrating to the ocean where they typically stay near the coastline and the mouths of larger rivers. In freshwater, found in small, low gradient streams and estuaries.	А	No suitable habitat present in BSA.
Steelhead, Klamath mountains province DPS	O. mykiss irideus	SSC	Spawns in gravel and small cobble substrates usually associated with riffle-and-run habitat types in coldwater streams.	А	No suitable habitat present in BSA.
Eulachon	Thaleichthys pacificus	FT	Spawns in lower reaches of rivers during peak spring flow events. Adults in the southern DPS are semelparous. Needs sand or coarse gravel for spawning substrate. Larvae are transported to estuaries and then to the ocean.	А	No suitable habitat present in BSA.
	Г	Π	Invertebrates	T	
Oregon silverspot butterfly	Speyeria zerene hippolyta	FT	Found in marine terrace and coastal headland meadows, stabilized dunes, and montane grasslands found on Mount Hebo and Fairview Mountain in Oregon (USFWS 2001). Requires early blue violet (<i>Viola adunca</i>), the larval host plant and nectar plants for adult butterflies.	А	No suitable habitat present in BSA; BSA south of known range.

Common Name	Scientific Name	Statusª	Habitat Requirements	Habitat Present/ Absent ^b	Rationale
Western bumble bee	Bombus occidentalis	SCE	Primarily found in shrubland and grassland. Typically nests in underground abandoned rodent burrows or other cavities.	А	No suitable habitat present in BSA.

Status:

Federal

FE = listed as endangered under the federal Endangered Species Act. listed as threatened under the federal Endangered Species Act.

FD = FPT = removed from federal Endangered Species Act list.

Federally proposed threatened.

State

ST = listed as threatened under the California Endangered Species Act.

SFP = designated as a fully protected species under the CFGC. State candidate endangered.

SCE = SCT = State candidate threatened.
SSC = State species of special concern.

Habitat Present/Absent:

= absent (no habitat present).

present (species is present).habitat present (habitat is or may be present). HP project footprint is in designated critical habitat.